



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*Foreign and insular statistical reports of countries and cities—Yearly and monthly.*

AUSTRALIA—*Queensland—Brisbane*.—Month of October, 1901. Estimated population, 119,428. Total number of deaths, 119, including diphtheria, 1; enteric fever, 3; whooping cough, 2; plague, 1, and 7 from tuberculosis.

BAHAMAS—*Governors Harbor*.—Three weeks ended January 4, 1902. Estimated population, 1,500. No deaths and no contagious diseases.

*Green Turtle Cay—Abaco*.—Two weeks ended January 1, 1902. Estimated population, 3,314. No deaths and no contagious diseases.

*Nassau*.—Two weeks ended January 4, 1902. Estimated population, 12,535. Number of deaths not reported. No contagious diseases reported.

BRITISH COLUMBIA—*Vancouver*.—Month of December, 1901. Estimated population, 26,000. Total number of deaths not reported. One death from scarlet fever reported.

CANADA—*Ontario—Hamilton*.—Month of December, 1901. Estimated population, 52,550. Total number of deaths, 60, including diphtheria, 3; enteric fever, 1, and 9 from tuberculosis.

ECUADOR—*Guayaquil*.—Two weeks ended November 9, 1901. Estimated population, 60,000. Total number of deaths, 344, including scarlet fever, 3, and 3 from smallpox.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 33 great towns in England and Wales during the week ended December 28, 1901, correspond to an annual rate of 21.1 per 1,000 of the aggregate population, which is estimated at 11,463,026. The highest rate was recorded in Manchester, viz, 29.3, and the lowest in Derby, viz, 14.3.

*Bradford*.—Two weeks ended December 28, 1901. Estimated population, 280,161. Total number of deaths, 200, including diphtheria, 1; enteric fever, 1; measles, 3; scarlet fever, 1, and 21 from tuberculosis.

*London*.—One thousand seven hundred and ninety-two deaths were registered during the week, including measles, 78; scarlet fever, 15; diphtheria, 28; whooping cough, 18; enteric fever, 6; smallpox, 25, and diarrhea and dysentery, 13. The deaths from all causes correspond to an annual rate of 20.6 per 1,000. In Greater London 2,427 deaths were registered, corresponding to an annual rate of 19.2 per 1,000 of the population. In the "outer ring" the deaths included 14 from diphtheria, 11 from measles, 7 from scarlet fever, 5 from smallpox, and 6 from whooping cough.

*Ireland*.—The average annual death rate represented by the deaths registered during the week ended December 28, 1901, in the 21 principal town districts of Ireland was 22.8 per 1,000 of the population, which is estimated at 1,079,708. The lowest rate was recorded in Tralee, viz, 0.0, and the highest in Limerick, viz, 34.2 per 1,000. In Dublin and suburbs 200 deaths were registered, including diphtheria, 2; enteric fever, 5, and 21 from tuberculosis.

*Scotland.*—The deaths registered in 8 principal towns during the week ended December 28, 1901, correspond to an annual rate of 22.6 per 1,000 of the population, which is estimated at 1,656,525. The lowest mortality was recorded in Leith, viz, 14.1, and the highest in Glasgow, viz, 24.8 per 1,000. The aggregate number of deaths registered from all causes was 720, including diphtheria, 8; measles, 36; scarlet fever, 7, and 7 from whooping cough.

ITALY—*Milan.*—Month of November, 1901. Estimated population, 496,156. Total number of deaths, 785, including diphtheria, 12; enteric fever, 19; measles, 2, and 78 from tuberculosis.

JAPAN—*Nagasaki.*—Ten days ended December 20, 1901. Estimated population, 131,700. Total number of deaths not reported. Three deaths from diphtheria and 3 from enteric fever reported.

PHILIPPINE ISLANDS—*Manila.*—Month of October, 1901. Population, 250,000. Total number of deaths, 885, including enteric fever, 6; measles, 1; plague, 7; smallpox, 1, and 68 from tuberculosis.

RUSSIA—*Riga.*—Month of September, 1901. Estimated population, 282,000. Total number of deaths, 588, including diphtheria, 6; enteric fever, 13; scarlet fever, 63; whooping cough, 17; smallpox, 4, and 50 from phthisis pulmonalis.

Month of October, 1901. Total number of deaths, 586, including diphtheria, 10; enteric fever, 12; measles, 1; scarlet fever, 66; whooping cough, 19; smallpox, 3, and 52 from phthisis pulmonalis.

ST. HELENA.—Four weeks ended November 16, 1901. Estimated population, 3,342. Total number of deaths, 7, including 1 from tuberculosis.